

Name: \_\_\_\_\_

AP Calculus AB - Definite Integral as the limit of a Riemann Sum  
By: Eiel Gonzalez, East Longmeadow HS (MA) and Brendan Murphy, John Bapst HS (ME)

Learning objective: Interpret the definite integral as the limit of a Riemann sum; interpret the limit of a Riemann sum as a definite integral.

Main calculus concept/formula:  $\lim_{n \rightarrow \infty} \sum_{k=1}^n f(a+k\Delta x) \Delta x = \int_a^b f(x) dx$

Part I: Translate the definite integral into a Riemann sum. Do NOT evaluate.

Example:  $\int_0^3 (x^2 - 1) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left[ \left( \left( \frac{3k}{n} \right)^2 - 1 \right) \left( \frac{3}{n} \right) \right]$

1)  $\int_1^2 (x+1) dx$

2)  $\int_2^4 (3x^2 - 1) dx$

3)  $\int_0^{\pi} (\sin x) dx$

4)  $\int_{-2}^1 (3x^2 + 2) dx$

5)  $\int_3^7 (2x^2 + 3x) dx$

1) Using the right and left approximation with 6 equal subdivisions each, estimate the distance traveled over  $[0, 3]$  if the velocity of an object at half-second intervals is:

t	0	0.5	1	1.5	2	2.5	3
v (ft/sec)	0	12	18	20	14	20	20

2) Find the distance of the object over the interval  $[0,3]$  in the previous problem using 3 subdivisions with a midpoint approximation.

3) Find the distance of the object over the interval  $[0, 3]$  in the problem 1 using 6 subdivisions with a trapezoidal approximation.

4) A rainstorm hit Portland, Maine, in October 1996, resulting in record rainfall. The rainfall rate  $R(t)$  on October 21 is recorded, in inches per hour, in the following table, where  $t$  is the number of hours since midnight. Compute the total rainfall during this 24-hour period using a right and a left approximation with 5 subdivisions.

t	0	3	5	10	16	24
R(t)	0.2	0.1	0.4	1.0	0.6	0.25

2015 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS

$t$ (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

3. Johanna jogs along a straight path. For  $0 \leq t \leq 40$ , Johanna's velocity is given by a differentiable function  $v$ . Selected values of  $v(t)$ , where  $t$  is measured in minutes and  $v(t)$  is measured in meters per minute, are given in the table above.

(a) Use the data in the table to estimate the value of  $v'(16)$ .

(b) Using correct units, explain the meaning of the definite integral  $\int_0^{40} |v(t)| dt$  in the context of the problem.

Approximate the value of  $\int_0^{40} |v(t)| dt$  using a right Riemann sum with the four subintervals indicated in the table.

(c) Bob is riding his bicycle along the same path. For  $0 \leq t \leq 10$ , Bob's velocity is modeled by  $B(t) = t^3 - 6t^2 + 300$ , where  $t$  is measured in minutes and  $B(t)$  is measured in meters per minute.

Find Bob's acceleration at time  $t = 5$ .

(d) Based on the model  $B$  from part (c), find Bob's average velocity during the interval  $0 \leq t \leq 10$ .

---

2016 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS

$t$ (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

1. Water is pumped into a tank at a rate modeled by  $W(t) = 2000e^{-t^2/20}$  liters per hour for  $0 \leq t \leq 8$ , where  $t$  is measured in hours. Water is removed from the tank at a rate modeled by  $R(t)$  liters per hour, where  $R$  is differentiable and decreasing on  $0 \leq t \leq 8$ . Selected values of  $R(t)$  are shown in the table above. At time  $t = 0$ , there are 50,000 liters of water in the tank.
- (a) Estimate  $R'(2)$ . Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.

2011 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS

$t$ (minutes)	0	2	5	9	10
$H(t)$ (degrees Celsius)	66	60	52	44	43

2. As a pot of tea cools, the temperature of the tea is modeled by a differentiable function  $H$  for  $0 \leq t \leq 10$ , where time  $t$  is measured in minutes and temperature  $H(t)$  is measured in degrees Celsius. Values of  $H(t)$  at selected values of time  $t$  are shown in the table above.

(a) Use the data in the table to approximate the rate at which the temperature of the tea is changing at time  $t = 3.5$ . Show the computations that lead to your answer.

(b) Using correct units, explain the meaning of  $\frac{1}{10} \int_0^{10} H(t) dt$  in the context of this problem. Use a trapezoidal sum with the four subintervals indicated by the table to estimate  $\frac{1}{10} \int_0^{10} H(t) dt$ .

(c) Evaluate  $\int_0^{10} H'(t) dt$ . Using correct units, explain the meaning of the expression in the context of this problem.

2017 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS

$h$ (feet)	0	2	5	10
$A(h)$ (square feet)	50.3	14.4	6.5	2.9

1. A tank has a height of 10 feet. The area of the horizontal cross section of the tank at height  $h$  feet is given by the function  $A$ , where  $A(h)$  is measured in square feet. The function  $A$  is continuous and decreases as  $h$  increases. Selected values for  $A(h)$  are given in the table above.
- (a) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the volume of the tank. Indicate units of measure.
- (b) Does the approximation in part (a) overestimate or underestimate the volume of the tank? Explain your reasoning.
- (c) The area, in square feet, of the horizontal cross section at height  $h$  feet is modeled by the function  $f$  given by  $f(h) = \frac{50.3}{e^{0.2h} + h}$ . Based on this model, find the volume of the tank. Indicate units of measure.